



Correction: Consensus statement on chronic pain treatment in cancer survivors

Keiko Mamiya¹ · Hiroki Iida² · Masako Iseki³ · Shigeki Yamaguchi⁴ · Hiroshi Yonekura⁵ · Hiroshi Ueno⁶ · Toshifumi Kosugi⁷ · Takeshi Sasara⁸ · Yumiko Takao⁹ · Toshifumi Takasusuki⁴ · Saori Hashiguchi¹⁰ · Naomi Hirakawa¹¹ · Yoko Sugiyama² · Keiko Yamada³ · Kenji Yamamoto¹²

Published online: 18 December 2024
© The Author(s) 2024

Correction: Journal of Anesthesia

<https://doi.org/10.1007/s00540-024-03427-0>

In the first sentence of the abstract, ‘the Japanese Society for Palliative Medicine (JSPM)’ should have been changed to ‘the Japan Society of Pain Clinicians (JSPC)’.

In the first sentence of the introduction, ‘the Japanese Society for Palliative Medicine (JSPM)’ should have been changed to ‘the Japan Society of Pain Clinicians (JSPC)’.

In the first sentence of the conclusion, ‘the Japanese Association of Pain Clinics (JAPC)’ should have been changed to ‘the Japan Society of Pain Clinicians (JSPC)’.

The original article has been corrected.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s00540-024-03427-0>.

✉ Keiko Mamiya
keiko@shinshu-u.ac.jp

- ¹ Division of Palliative Medicine, Shinshu Cancer Center, Shinshu University Hospital, 3-1-1 Asahi, Matsumoto 390-8621, Japan
- ² Gifu University/Anesthesiology and Pain Relief Center, Central Japan International Medical Center, Minokamo, Japan
- ³ Department of Anesthesiology and Pain Medicine, Faculty of Medicine, Juntendo University, Bunkyo, Japan
- ⁴ Department of Anesthesiology, Dokkyo Medical University School of Medicine, Mibu, Japan
- ⁵ Department of Anesthesiology and Pain Medicine, Fujita Health University Bantane Hospital, Nagoya, Japan

- ⁶ Department of Anesthesiology, Kyoto Prefectural University of Medicine, Kyoto, Japan
- ⁷ Department of Palliative Care, Saga-Ken Medical Center Koseikan, Saga, Japan
- ⁸ Yuuikai Tomishiro Central Hospital, Total Pain Center, Tomigusuku, Japan
- ⁹ Department of Pain Medicine, Hyogo Medical University Hospital, Nishinomiya, Japan
- ¹⁰ Department of Palliative Medicine, St. Marianna University School of Medicine, Kawasaki, Japan
- ¹¹ Department of Anesthesiology and Pain Clinic, Hirakawa Hospital, Tokyo, Japan
- ¹² Department of Palliative Care, Hokkaido Cancer Center, Hokkaido, Japan